

## Chapter 2.

# Soil and Fertilizer Management for Vegetable Production in Florida

E.H. Simonne and G.J. Hochmuth

### BEST MANAGEMENT PRACTICES

With the passage of the Federal Clean Water Act (FCWA) in 1972, states were required to assess the impacts of non-point sources of pollution on surface and ground waters, and establish programs to minimize them. Section 303(d) of the FWCA also requires states to identify impaired water bodies and establish total maximum daily loads (TMDLs) for pollutants entering these water bodies. Water quality parameters targeted by the TMDLs and involving vegetable production are concentrations of nitrate, phosphate, and total dissolved solids these waters. A TMDL establishes the maximum amount of pollutant a water body can receive and still keep its water quality parameters consistent with its intended use (swimming, fishing, or potable uses). The establishment of the TMDLs is currently underway and they will be implemented through a combination of regulatory, non-regulatory, and incentive-based measures. Best Management Practices (BMPs) are specific cultural practices aimed at reducing the load of a specific compound, while maintaining or increasing economical yields. They are tools available to vegetable growers to achieve the TMDLs. BMPs are intended to be educational, economically sound, environmentally effective, and based on science. It is important to recognize that BMPs do not aim at becoming an obstacle to vegetable production. Instead, they should be viewed as a means to balance economical vegetable production with environmental responsibility.

The BMPs that will apply to vegetable production in Florida are described in the 'Agronomic and Vegetable Crop Water Quality/Water Quantity BMP Manual for Florida'. This 190-page manual has been developed between 2000 and 2002 through a cooperative effort between state agencies, water management districts and commodity groups, and under the scientific leadership of the University of Florida's Institute of Food and Agricultural Sciences (UF/IFAS). The manual has undergone a thorough scientific review at the beginning of 2003. In the next step, the manual will be presented to stakeholders and state commodity groups for feed back. The latest manual draft may be consulted on-line at <http://www.floridaagwaterpolicy.com/PDFs/BMPs/vegetable&agronomicCrops.pdf>. BMPs are 1-to-3 page long chapters that include a picture, a working definition of the topic, list specific

things to do (BMPs) as well as things to avoid (pitfalls), and present existing applicable technical criteria together with additional references.

The vegetable BMPs have adopted all current IFAS recommendations, including those for fertilizer and irrigation management. Through the implementation of a series of targeted cultural practices (the BMPs), growers should be able to reconcile economical profitability and responsible use of water and fertilizer. At the field level, adequate fertilizer rates should be used together with irrigation scheduling techniques and crop nutritional status monitoring tools (leaf analysis, petiole sap testing). In the BMP manual, adequate fertilizer rates may be achieved by combinations of IFAS recommended rates and supplemental fertilizer applications.

### SOILS

Vegetables are grown on more than 300,000 acres in various soil types throughout the state. These soil types include sandy soils, sandy loam soils, Histosols (organic muck), and calcareous marl soils. Each soil group is described below.

#### Sands

Sandy soils make up the dominant soil type for vegetable production in Florida (Fig. 2-1). Vegetables are produced on sandy soils throughout the Florida peninsula and on sandy soils and sandy loams in the panhandle. Sandy soils have the advantage of ease of tillage and they can produce the earliest vegetable crops for a particular region. Sandy soils allow timely production operations such as planting and harvesting. Sandy soils, however, have the disadvantage that mobile nutrients such as nitrogen and potassium can be leached by heavy rain or over irrigation. Therefore, sands must be managed carefully with regard to fertility programs. Sands hold very little water; therefore, irrigation management is more critical compared to other soil types used for vegetable production in Florida. Nearly all vegetable crops produced in Florida can be successfully grown on sandy soils. The major vegetable crops such as tomatoes, peppers, potatoes, watermelons, strawberries, and cabbage are grown commonly on sandy soils.

**Table 1.** Fertilizer nutrients required by plants.

Nutrient	Deficiency symptoms	Occurrence
Nitrogen (N)	Stems thin, erect, hard. Leaves small, yellow; on some crops (tomatoes) undersides are reddish. Lower leaves affected first.	On sandy soils especially after heavy rain or after overirrigation. Also on organic soils during cool growing seasons.
Phosphorus (P)	Stems thin and shortened. Leaves develop purple color. Older leaves affected first. Plants stunted and maturity delayed.	On acidic soils or very alkaline soils. Also when soils are cool and wet.
Potassium (K)	Older leaves develop gray or tan areas on leaf margins. Eventually a scorch appears on the entire margin.	On sandy soils following leaching rains or overirrigation.
Boron (B)	Growing tips die and leaves are distorted. Specific diseases caused by boron deficiency include brown curd and hollow stem of cauliflower, cracked stem of celery, blackheart of beet, and internal browning of turnip.	On soils with pH above 6.8 or on sandy, leached soils, or on crops with very high demand such as cole crops.
Calcium (Ca)	Growing-point growth restricted on shoots and roots. Specific deficiencies include blossom-end rot of tomato, pepper and watermelon, brownheart of escarole, celery blackheart, and cauliflower or cabbage tipburn.	On strongly acidic soils, or during severe droughts.
Copper (Cu)	Yellowing of leaves, stunting of plants. Onion bulbs are soft with thin, pale scales.	On organic soils or occasionally new mineral soils.
Iron (Fe)	Distinct yellow or white areas between veins on youngest leaves.	On soils with pH above 6.8.
Magnesium (Mg)	Initially older leaves show yellowing between veins, followed by yellowing of young leaves. Older leaves soon fall.	On strongly acidic soils, or on leached sandy soils.
Manganese (Mn)	Yellow mottled areas between veins on youngest leaves, not as intense as iron deficiency.	On soils with pH above 6.4.
Molybdenum (Mo)	Pale, distorted, narrow leaves with some interveinal yellowing of older leaves, e.g. whiptail disease of cauliflower.	On very acidic soils.
Zinc (Zn)	Small reddish spots on cotyledon leaves of beans; light areas (white bud) of corn leaves.	On wet, cold soils in early spring or where excessive phosphorus is present.
Sulfur (S)	General yellowing of younger leaves and reduced growth.	On very sandy soils, low in organic matter, especially following continued use of sulfur-free fertilizers and especially in areas that receive little atmospheric sulfur.
Chlorine (Cl)	Deficiencies very rare.	Usually only under laboratory conditions.

### Histosols

Histosols are organic soils which occur in areas throughout the peninsula, especially in southern and central Florida (Fig. 2-2). Large organic deposits used for vegetable production occur south of Lake Okeechobee. Smaller pockets of “muck” occur throughout central and northern Florida. Histosols consist largely of decomposing plant material and are largely underlain by calcareous deposits. Muck soils have large water and nutrient holding capacities and are used to produce crops such as the leafy vegetables (leaf lettuce, and various greens), celery, sweet corn, and radishes. With time, the organic matter decomposes and the muck subsides. Thus, the pH in the muck can increase because of proximity to the underlying calcareous material. Muck subsidence causes problems for water and nutrient management. The increase in pH due to subsidence and also to the practice of flooding the Histosols to reduce oxidation can result in increased requirements of phosphorus and micronutrients. These

nutrients can be fixed by the high pH of the soil. Nutrient management in these situations should involve banding than increased rates of nutrients.

### Calcareous Rock and Marl

The calcareous soils in southern Florida (Miami-Dade County) consist of two phases, rockland and marl. Rockland soils are calcium carbonate soils consisting of particles that range from sand-like in size to pebble and gravel (Fig. 2-3). The rockland soils are extremely shallow, about 4 to 6 inches deep. The marl is the fine-textured, clay-like phase of the calcium carbonate soils. Tomatoes, beans, summer squash, okra, sweet corn, boniato, and strawberries can be produced in the winter months on the rockland soils of Dade County. Potatoes, malanga, and sweet corn are produced on the marl. Both soils have extremely high pH, therefore, nutrients such as phosphorus and micronutrients must be banded to ensure availability.

**Table 2.** Mehlich-1 (double-acid) interpretations for vegetable crops in Florida.

Element	Very low	Low	Medium	High	Very high
P	<10	10-15	16-30	31-60	>60
K	<20	20-35	36-60	61-125	>125
Mg	<10	10-20	21-40	41-60	>60
Ca	<100	100-200	201-300	301-400	>400

**Table 3.** Interpretations of Mehlich-1 soil tests for micronutrients.

	Soil pH (mineral soils only)		
	5.5 - 5.9	6.0 - 6.4	6.5 - 7.0
	parts per million		
Test level below which there may be a crop response to applied copper.	0.1 - 0.3	0.3 - 0.5	0.5
Test level above which copper toxicity may occur.	2.0 - 3.0	3.0 - 5.0	5.0
Test level below which there may be a crop response to applied manganese.	3.0 - 5.0	5.0 - 7.0	7.0 - 9.0
Test level below which there may be a crop response to applied zinc.	0.5	0.5 - 1.0	1.0 - 3.0

### SOIL TESTING

Plants require 16 elements for normal growth and reproduction (Table 1). The crop nutrient requirement (CNR) for a particular element is defined as the total amount in lb/A of that element needed by the crop to produce economic optimum yield. This concept of economic optimum yields is important for vegetables because a certain amount of nutrients might produce a moderate amount of biomass, but produce negligible marketable product due to small fruit size. Fruit size and quality must be considered in the crop nutrient requirement concept for vegetables.

The crop nutrient requirement can be satisfied from many sources, including soil, water, air, organic matter, or fertilizer. For example, the CNR of potassium (K) can be supplied from K-containing minerals in the soil, from K retained by soil organic matter, or from K fertilizers.

The CNR for a crop is determined from field experiments that test the yield response to levels of added fertilizer. For example, a watermelon study involving K might be conducted on a soil which tests very low in extractable K. In this situation, the soil can be expected to contribute only a small amount of K for optimum watermelon growth and yield, and K must be supplied largely from fertilizer. The researcher plots the relationship between crop yield and fertilizer rate. The CNR is equivalent to the fertilizer rate above which no significant increases in yield are expected. The CNR values derived from such experiments take into account factors such as fertilizer efficiencies of the soils. These efficiencies include fertilizer leaching or fertilizer nutrient fixing capability of the soil. If data are available from several experiments, then reliable estimates of CNR values can be made. Using the CNR concept

when developing a fertilizer program will ensure optimum, economic yields while minimizing both pollution from overfertilization and loss of yield due to underfertilization.

The CNR values are those amounts of nutrients needed to produce optimum, economic yields from a fertilization standpoint. It is important to remember that these nutrient amounts are supplied to the crop from both the soil and the fertilizer. The amounts are applied as fertilizers only when a properly calibrated soil test indicates very small extractable amounts of these nutrients to be present in the soil. Therefore, soil testing must be conducted to determine the exact contribution from the soil to the overall CNR. Based on such tests, the amount of fertilizer that is needed to supplement the nutrition component of the native soil can be calculated (Table 2 and 3).

It is important that soil samples represent the field or management unit to be fertilized. A competent soil testing laboratory that uses calibrated methodologies should analyze the samples. Not all laboratories can provide accurate fertilizer recommendations for Florida soils.

### LIMING

Current University of Florida standardized recommendations call for maintaining soil pH between 6.0 and 6.5 (Table 4). However, some vegetables, such as watermelon, will perform normally at lower soil pH as long as large amounts of micronutrients are not present in the soil. A common problem in Florida has been overliming, resulting in high soil pH. Overliming and resulting high soil pH can tie up micronutrients and restrict their availability to the

**Table 4.** A general guideline to crop tolerance of mineral soil acidity.<sup>1</sup>

Slightly tolerant (pH 6.8-6.0)		Moderately tolerant (pH 6.8-5.5)		Very tolerant (pH 6.8-5.0)
Beet	Leek	Bean, snap	Mustard	Endive
Broccoli	Lettuce	Bean, lima	Pea	Potato
Cabbage	Muskmelon	Brussels sprouts	Pepper	Shallot
Cauliflower	Okra	Carrot	Pumpkin	Sweetpotato
Celery	Onion	Collard	Radish	Watermelon
Chard	Spinach	Corn	Squash	
		Cucumber	Tomato	
		Eggplant	Turnip	
		Kale		

<sup>1</sup> From Donald N. Maynard and George J. Hochmuth, Knott's Handbook For Vegetable Growers, 4th edition (1997). Reprinted by permission of John Wiley & Sons, Inc.

**Table 5.** Liming materials.

Material	Formula	Amount of Material to be used to equal	
		1 ton of Calcium Carbonate <sup>1</sup>	Neutralizing value <sup>2</sup> (%)
Calcium carbonate, calcite, hi-cal lime	CaCO <sub>3</sub>	2,000lbs	100
Calcium-magnesium carbonate, dolomite	CaCO <sub>3</sub> • MgCO <sub>3</sub>	1,850lbs	109
Calcium oxide, burnt lime	CaO	1,100lbs	179
Calcium hydroxide, hydrated lime	Ca(OH) <sub>2</sub>	1,500lbs	136
Calcium silicate, slag	CaSiO <sub>3</sub>	2,350lbs	86
Magnesium carbonate	MgCO <sub>3</sub>	1,680lbs	119

<sup>1</sup> Calculated as (2000 x 100) / neutralizing value (%).  
<sup>2</sup> The higher the neutralizing value, the greater the amount of acidity that is neutralized per unit weight of material.

crop. Overliming also can reduce the accuracy with which a soil test can predict the fertilizer component of the CNR.

It is important, however, not to allow soil pH to drop below approximately 5.5 for most vegetable production, especially where micronutrient levels in the soil may be high due to a history of micronutrient fertilizer and micronutrient-containing pesticide applications. When soil pH decreases in such soils, the solubility of micronutrients can increase to levels that may become toxic to plants.

Irrigation water from wells in limestone aquifers is an additional source of liming material usually not considered in many liming programs. The combination of routine additions of lime and use of alkaline irrigation water has resulted in soil pH greater than 8.0 for many sandy soils in south Florida. To measure the liming effect of irrigation, have a water sample analyzed for total bicarbonates and carbonates annually, and the results converted to pounds of calcium carbonate per acre. Include this information in your decisions concerning lime.

It should be evident that liming (Table 5), fertilization (Table 6), and irrigation programs are closely related to each other. An adjustment in one program will often influence the other. To maximize overall production efficiency, soil and water testing must be made a part of any fertilizer management program.

## MANURES/SLUDGE

Waste organic products, including animal manures, composted organic matter, and sludges, contain nutrients (Table 7) that can enhance plant growth. These materials decompose when applied to the soil, releasing nutrients that vegetable crops can absorb and utilize in plant growth. The key to proper use of organic materials as fertilizers comes in the knowledge of the nutrient content and the decomposition rate of the material. Many laboratories offer organic material analyses to determine specific nutrient contents. Growers contemplating using organic materials as fertilizers should have an analysis of the material before determining the rate of application. In the case of materials such as sludges, it is important to have knowledge about the type of sludge to be used. Certain classes of sludge are not appropriate for vegetable production, and in fact may not be permitted for land application. Decomposition rates of organic materials in warm sandy soils in Florida are rapid. Therefore, there will be relatively small amounts of residual nutrients remaining for succeeding crops. Organic materials are generally similar to mixed chemical fertilizers in that the organic waste supplies an array of nutrients, some of which may not be required on a particular soil. For example, the P in poultry manure would not be required on a soil already testing high in phosphate. Usually application rates of organic wastes are determined largely by the N content. Organic waste materials can con-

**Table 6.** Effect of some fertilizer materials on soil pH.

Fertilizer material	Approximate calcium carbonate equivalent (lb) <sup>1</sup>
Ammonium nitrate	-1200
Ammonium sulfate	-2200
Anhydrous ammonia	-3000
Diammonium phosphate	-1250 to -1550
Potassium chloride	0
Sodium-potassium nitrate	+550
Nitrogen solutions	-759 to -1800
Normal (ordinary) superphosphate	0
Potassium nitrate	+520
Potassium sulfate	0
Potassium-magnesium sulfate	0
Triple (concentrated) superphosphate	0
Urea	-1700

<sup>1</sup>A minus sign indicates the number of pounds of calcium carbonate needed to neutralize the acid formed when one ton of fertilizer is added to the soil.

**Table 7.** Average nutrient concentration of selected organic fertilizers.

Product	N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O
	% dry weight		
Blood	13	2	1
Activated sewage	6	3	0.2
Fish meal	10	6	0
Bone meal	3	22	0
Cotton seed meal	6	3	1.5
Peanut meal	7	1.5	1.2
Soybean meal	7	1.2	1.5
Dried commercial manure products			
Stockyard	1	1	2
Cattle	2	3	3
Chicken	1.5	1.5	2

tribute to groundwater or surface water pollution if applied in rates in excess of the crop nutrient requirement for a particular vegetable crop. Therefore, it is important to understand the nutrient content and the decomposition rate of the organic waste material, and the P-holding capacity of the soil.

## N, P, K, NUTRIENT SOURCES

Nitrogen can be supplied in both nitrate and ammoniacal forms (Table 8). Nitrate-nitrogen is generally the preferred form for plant uptake in most situations, but ammoniacal N can be absorbed directly or after conversion to nitrate-N by soil microbes. Since this rate of conversion is reduced in cold, fumigated, or strongly acidic soils, it is recommended that under such conditions 25%

to 50% of the N be supplied from nitrate sources. This ratio is not as critical for unfumigated or warm soils. For more information on nutrient sources, consult Florida Extension Circular 225-E, *Commercial Vegetable Fertilization Guide*.

Phosphorus (P) can be supplied from several sources, including normal and triple superphosphate, diammonium phosphate, monopotassium phosphate, and monoammonium phosphate. All sources can be effective for plant nutrition on sandy soil. However, on soils that test very low in native micronutrient levels, diammonium phosphate in mixtures containing micronutrients reduces yields when banded in large amounts. Availability of P also can be reduced with use of diammonium phosphate compared to use of triple superphosphate. Negative effects of diammonium phosphate can be eliminated by using it for only a portion of the P requirement and by broadcasting this material in the bed.

Potassium (K) can also be supplied from several sources, including potassium chloride, potassium sulfate, potassium nitrate, and potassium-magnesium sulfate. If soil-test-predicted amounts of K fertilizer are adhered to, there should be no concern about the K source or its relative salt index.

## Ca, S, AND Mg

The secondary nutrients calcium (Ca), sulfur (S), and magnesium (Mg) have not been a common problem in Florida. Calcium usually occurs in adequate supply for most vegetables when the soil is limed. If the Mehlich-1 soil Ca index is above 300 ppm, it is unlikely that there will be a response to added Ca. Maintaining correct moisture levels in the soil by irrigation will aid in Ca supply to the roots. Calcium is not mobile in the plant; therefore, foliar sprays of Ca are not likely to correct deficiencies. It is difficult to place enough foliar-applied Ca at the growing point of the plant on a timely basis.

Sulfur deficiencies have seldom been documented for Florida vegetables. Sulfur deficiency would most likely occur on deep, sandy soils low in organic matter after leaching rains. If S deficiency has been diagnosed, it can be corrected by using S-containing fertilizers such as magnesium sulfate, ammonium sulfate, potassium sulfate, normal superphosphate, or potassium-magnesium sulfate. Using one of these materials in the fertilizer blend at levels sufficient to supply 30 to 40 lb S/A should prevent S deficiencies. In many areas of Florida, ample S will be supplied to vegetable crops through the irrigation water because sulfur can be present in the aquifer.

Magnesium deficiency may be a problem for vegetable production; however, when the Mehlich-1 soil-test index

**Table 8.** Some commonly used fertilizer sources.

<b>Nutrient</b>	<b>Fertilizer source</b>	<b>Nutrient content (%)</b>
Nitrogen (N)	Ammonium nitrate	34
	Ammonium sulfate	21
	Calcium nitrate	15.5
	Diammonium phosphate	18
	Potassium nitrate (nitrate of potash)	13
	Urea	46
	Sodium-potassium nitrate (nitrate of soda-potash)	13
Phosphorus (P <sub>2</sub> O <sub>5</sub> )	Normal (ordinary) superphosphate	20
	Triple (concentrated) superphosphate	46
	Diammonium phosphate	46
	Monopotassium phosphate	53
Potassium (K <sub>2</sub> O)	Potassium chloride (muriate of potash)	60
	Potassium nitrate	44
	Potassium sulfate (sulfate of potash)	50
	Potassium-magnesium sulfate (sulfate of potash-magnesia)	22
	Sodium-potassium nitrate	14
	Monopotassium phosphate	34
Calcium (Ca)	Calcic limestone	32
	Dolomite	22
	Gypsum	23
	Calcium nitrate	19
	Normal superphosphate	20
	Triple superphosphate	14
Magnesium (Mg)	Dolomite	11
	Magnesium sulfate	10
	Magnesium oxide	55
	Potassium-magnesium sulfate	11
Sulfur (S)	Elemental sulfur	97
	Ammonium sulfate	24
	Gypsum	18
	Normal superphosphate	12
	Magnesium sulfate	14
	Potassium-magnesium sulfate	22
	Potassium sulfate	18
Boron (B)	Borax	11
	Fertibor <sup>1</sup>	14.9
	Granubor <sup>1</sup>	14.3
	Solubor <sup>1</sup>	20.5
Copper (Cu)	Copper sulfate, monohydrate	35
	Copper sulfate, pentahydrate	25
	Cupric oxide	75
	Cuprous oxide	89
	Copper chloride	17
	Chelates (CuEDTA)	13
	(CuHEDTA)	6
Iron (Fe)	Ferrous sulfate	20
	Ferric sulfate	20
	Chelates (FeHEDTA)	5 to 12
Manganese (Mn)	Manganous sulfate	28
	Manganous oxide	68
	Chelates (MnEDTA)	5 to 12
Molybdenum (Mo)	Ammonium molybdate	54
	Sodium molybdate	39
Zinc (Zn)	Zinc sulfate	36
	Zinc oxide	80
	Zinc chloride	50
	Chelates (ZnEDTA)	6 to 14
	(ZnHEDTA)	6 to 10

<sup>1</sup>Mention of a trade name does not imply a recommendation over similar materials.

**Table 9.** Recommendations for foliar applications of plant nutrients.

Nutrient	Source	Foliar application (lb product per acre)
Boron	Borax	2 to 5
	Solubor <sup>1</sup>	1 to 1.5
Copper	Copper sulfate	2 to 5
Iron	Ferrous sulfate	2 to 3
	Chelated iron	0.75 to 1
Manganese	Manganous sulfate	2 to 4
Molybdenum	Sodium molybdate	0.25 to 0.50
Zinc	Zinc sulfate	2 to 4
	Chelated zinc	0.75 to 1
Calcium	Calcium chloride	5 to 10
	Calcium nitrate	5 to 10
Magnesium	Magnesium sulfate	10 to 15

<sup>1</sup> Mention of a trade name does not imply a recommendation over similar materials.

for Mg is below 15 ppm, 30 lb Mg/A will satisfy the Mg CNR. If lime is also needed, Mg can be added by using dolomite as the liming material. If no lime is needed, then the Mg requirement can be satisfied through use of magnesium sulfate or potassium-magnesium sulfate. Blending of the Mg source with other fertilizer(s) to be applied to the soil is an excellent way of ensuring uniform application of Mg to the soil.

### MICRONUTRIENTS

It has been common in Florida vegetable production to routinely apply a micronutrient package. This practice has been justified on the basis that these nutrients were inexpensive and their application appeared to be insurance for high yields. In addition, there were few research data and a lack of soil-test calibrations to guide judicious application of micronutrient fertilizers. Compounding the problem has been the vegetable industry's use of micronutrient-containing pesticides for disease control. Copper (Cu), manganese (Mn), and zinc (Zn) from pesticides have tended to accumulate in the soil.

This situation has forced some vegetable producers to overlime in an effort to avoid micronutrient toxicities. Data have now been accumulated which permit a more accurate assessment of micronutrient requirements (Table 3). Growers are encouraged to have a calibrated micronutrient soil test conducted and to refrain from shotgun micronutrient fertilizer applications. It is unlikely that micronutrient fertilizers will be needed on old vegetable land, especially where micronutrients are being applied regularly via recommended pesticides. A micronutrient soil test every 2 to 3 years will provide recommendations for micronutrient levels for crop production.

### FOLIAR FERTILIZATION

Foliar fertilization should be thought of as a last resort for correcting a nutrient deficiency (Table 9). The plant leaf is structured in such a way that it naturally resists easy infiltration by fertilizer salts. Foliar fertilization most appropriately applies to micronutrients and not to macronutrients such as N, P, and K. Foliar applications of N, P, and/or K are not needed where proper soil-directed fertilizer programs are in use. Leaves cannot absorb sufficient nutrients (without burning the leaves) to correct any deficiency. Some benefit from macronutrient foliar sprays probably results when nutrients are washed by rain or irrigation water off the leaf surface into the soil. The nutrient then may enter the plant via the roots. Amounts of macronutrients recommended on the label of most commercial foliar products are so minuscule compared to nutrition derived from the soil that benefit to the plant is highly unlikely. Additionally, fertilizer should only be added if additional yield results, and research with foliar-nutrient applications has not clearly documented a yield increase for vegetables.

In certain situations, temporary deficiencies of Mn, Fe, Cu, or Zn can be corrected by foliar application. Examples include vegetable production in winter months when soils are cool and roots cannot extract adequate amounts of micronutrients, and in cases where high pH (marl and Rockdale soils) fixes broadcast micronutrients into unavailable forms. Micronutrients are so termed because small, or micro, amounts are required to satisfy the CNR. Such micro amounts may be supplied adequately through foliar applications to correct a temporary deficiency.

Boron is highly immobile in the plant. To correct boron deficiencies, small amounts of boron must be applied frequently to the young tissue or buds.

Any micronutrient should be applied **only** when a specific deficiency has been clearly diagnosed. Do not make unneeded applications of micronutrients. There is a fine line between adequate and toxic amounts of these nutrients. Indiscriminate application of micronutrients may reduce plant growth and restrict yields because of toxicity. Compounding the problem is the fact that the micronutrients can accumulate in the soil to levels which may threaten crop production on that soil. An important part of any micronutrient program involves careful calculations of all micronutrients being applied, from all sources.

### LIQUID VS. DRY FERTILIZER

There is no difference in response of crops to similar amounts of nutrients when applied in either liquid or dry form. Certain situations (use of drip irrigation or injection

wheel) require clear or true solutions. However, sidedress applications of fertilizer can be made equally well with dry or liquid forms of nutrients.

The decision to use liquid or dry fertilizer sources should depend largely on economics and on the type of application equipment available. The cost per unit of nutrient (e.g., dollars per unit of actual N) and the combination of nutrients provided should be used in any decision-making process.

## CONTROLLED-RELEASE FERTILIZERS (CRF)

Several brands of controlled-release fertilizers are available for supplying N. Some vegetables increase in yield when controlled-release fertilizers, such as polymer-coated or sulfur-coated urea, or isobutylidene-diurea, are used to supply a portion of the N requirement. Although more expensive, these materials may be useful in reducing fertilizer losses through leaching, in decreasing soluble salt damage, and in supplying adequate fertilizer for long-term crops such as strawberry or pepper. Controlled-release potassium fertilizers also have been demonstrated to be beneficial for several vegetables. It is essential to match the nutrient release pattern of the CRF with the crop's uptake pattern.

## SOLUBLE SALTS

Overfertilization or placement of fertilizer too close to the seed or root leads to soluble salt injury or "fertilizer burn." Fertilizer sources differ in their capacity to cause soluble salt injury. Therefore, where there is a history of soluble salt problems, or where irrigation water is high in soluble salts, choose low-salt index fertilizer sources, and broadcast or split-apply the fertilizer.

## STARTER FERTILIZER

A true starter fertilizer is a soluble fertilizer, generally high in P, used for establishment of young seedlings and transplants. Starter fertilizers generally work best if a small amount of N and K is present along with the P. Starters represent a very small percentage of the overall fertilizer amount but are very important in establishing crops in cool, damp soils. They can be applied with the planter at 2 inches to the side of the seed and 2 inches deep or can be dissolved in the transplant water and applied in the furrow.

## FERTILIZER PLACEMENT

Fertilizer rate and placement must be considered together. Banding low amounts of fertilizer too close to plants

can result in the same amount of damage as broadcasting excessive amounts of fertilizer in the bed.

Because P movement in most soils is minimal, it should be placed in the root zone. Banding is generally considered to provide more efficient utilization of P by plants than broadcasting. This is especially true on the high P-fixing calcareous soils. Where only small amounts of fertilizer P are to be used, it is best to band. If broadcasting P, a small additional amount of starter P near the seed or transplant may improve early growth, especially in cool soils. For wide-row crops such as watermelons, the modified broadcast method where fertilizer is broadcast only in the bed area provides more efficient use of fertilizer than complete broadcasting.

Micronutrients can be broadcast with the P and incorporated in the bed area. On the calcareous soils, micronutrients, such as Fe, Mn, and B, should be banded or applied foliarly.

Since N and, to a lesser extent, K are mobile in sandy soils, they must be managed properly to maximize crop uptake. Plastic mulch helps retain these nutrients in the soil. Under non-mulched systems, split applications of these nutrients must be used to reduce losses to leaching. Here, up to one-half of the N and K may be applied to the soil at planting or shortly after that time. The remaining fertilizer is applied in one or two applications during the early part of the growing season. Splitting the fertilizer applications also will help reduce the potential for soluble salt damage to the plants.

When using plastic mulch, fertilizer placement depends on the type of irrigation system (seep or overhead) and on whether drip tubing or the liquid fertilizer injection wheel are to be used.

With overhead irrigation, all fertilizer can be incorporated in the bed before mulch application. However, when large amounts of fertilizer are required, such as on virgin soil, a portion of the fertilizer can be banded to reduce salt damage. The fertilizer for crops grown on Rockdale soils can be banded, although research has shown equal yields by some crops with incorporation in the bed.

With seep irrigation, all P and micronutrients should be incorporated in the bed. Apply 10 to 20% (but not more) of the N and K with the P. The remaining N and K should be placed in narrow bands on the bed shoulders, the number of which depend on the crop and number of rows per bed. These bands should be placed in shallow (2- to 2½-inch-deep) grooves. This placement requires that adequate bed moisture be maintained so that capillarity is not broken. Otherwise, fertilizer will not move to the root zone.



Excess moisture can result in fertilizer leaching. Fertilizer and water management programs are linked. Maximum fertilizer efficiency is achieved only with close attention to water management.

Under either system above, fertilizing with drip irrigation or with a liquid fertilizer injection wheel might be suitable alternatives to the placement of all N and K in or on the bed prior to mulching.

In cases where supplemental sidedressing of mulched crops is needed, applications of liquid fertilizer can be made through the mulch with a liquid fertilizer injection wheel (Fig. 2-4). This implement is mounted on a tool bar and, using 30 to 40 psi pressure, injects fertilizer through a hole pierced in the mulch.

## SUPPLEMENTAL FERTILIZER AND BMPS

The use of supplemental fertilizer depends mainly on the amount of rainfall and the length of the season. The need for supplemental fertilizer will be minimized when the soil-test-predicted fertilizer requirements are properly managed to reduce losses to leaching. In most cases, N and K are of most concern because of leaching, although sidedressing of P might be needed on the phosphate-fixing calcareous soils, especially during cool periods. If required, supplemental applications of N and K should be made as a band or swath just ahead of the advancing root system. Place the fertilizer deep enough in the soil to be in contact with moisture. In many cases, liquid fertilizers are easiest to use and can be easily knifed into the soil. Sidedress applications must only be made when there is no danger of shoot or root damage from the fertilizer application machinery.

## SUPPLEMENTAL FERTILIZER APPLICATIONS

In practice, supplemental fertilizer applications allow vegetable growers to numerically apply fertilizer rates higher than the standard IFAS recommended rates when growing conditions require to do so. The two main growing conditions that may require supplemental fertilizer applications are leaching rains and extended harvest periods. Applying additional fertilizer under the following three circumstances is part of the current IFAS fertilizer recommendations. Supplemental N and K fertilizer applications may be made under three circumstances:

1. For vegetable crops grown on bare ground with seepage irrigation and without drip irrigation, a 30 lbs/acre of N and /or 20 lbs/acre of K<sub>2</sub>O supplemental application is allowed after a leaching rain. A leaching rain occurs when it rains at least 3 inches in 3 days, or 4 inches in 7 days.

2. For all vegetable crops grown on any production system with one of the IFAS recommended irrigation scheduling methods, a supplemental fertilizer application is allowed when nutrient levels in the leaf or in the petiole fall below the sufficiency ranges. For bare ground production, the supplemental amount allowed is 30 lbs/acre of N and/or 20 lbs/acre of K<sub>2</sub>O. For drip irrigated crops, the supplemental amount allowed is 1.5 to 2.0 lbs/A/day for N and/or K<sub>2</sub>O for one week.
3. Supplemental fertilizer applications are allowed when, for economical reasons, the harvest period has to be longer than the typical harvest period. When the results of tissue analysis and/or petiole testing are below the sufficiency ranges, a supplemental 30 lbs/acre N and/or 20 lbs/acre of K<sub>2</sub>O may be made for each additional harvest for bare ground production. For drip-irrigated crops, the supplemental fertilizer application is 1.5 to 2.0 lbs/A/day for N and/or K<sub>2</sub>O until the next harvest. A new leaf analysis and/or petiole analysis is required to document the need for additional fertilizer application for each additional harvest.

## DOUBLE-CROPPING

Successive cropping of existing mulched beds is a good practice in order to make effective use of the polyethylene mulch and fumigant (Fig. 2-5). Double-cropping also can make use of residual fertilizer in the beds. If fertilizer-N applications and amounts were properly managed for the first crop, then there should be negligible amounts of N-fertilizer remaining in the beds. The practice of adding extra fertilizer to the beds when planting the first crop, thinking that this fertilizer will aid growth of the second crop is strongly discouraged. The extra fertilizer could contribute to soluble-salt damage to the first crop, and might still be leached from the root zone before the second crop is established.

A drip-irrigation system can be used to supply adequate nutrition to each crop in a double crop system. In most cases, only N and K may be needed for the second crop. Amounts of P and micronutrients (if any) used for the first crop will likely remain adequate for the second crop as well. Soil testing of a sample taken from the bed away from any fertilizer bands will help determine P or micronutrient needs, assuming that these nutrients were broadcast in the bed prior to planting the first crop.

If N for the first crop was not applied in excess of the CNR, then the second crop should receive an amount of N equal to its own CNR. Potassium requirements of the second crop can be determined as for P in cases where the K for the first crop was incorporated in the bed. Potassium

requirements for the second crop are more difficult to determine in cases where K for the first crop was banded. A moderate amount of residual K will probably remain in the bed from the application to the first crop. Therefore, K requirements for the second crop will likely be slightly less than the CNR value for the chosen crop.

Once the crop fertilizer requirements have been ascertained, the needed nutrition may be applied through the drip system. Where drip irrigation is not being used, a liquid injection wheel can be used to place fertilizer in the bed for the second crop.

### LINEAR BED FOOT (LBF) SYSTEM FOR FERTILIZER APPLICATION

The University of Florida Extension Soil Testing Laboratory (ESTL) employs the Standardized Fertilizer Recommendation System in which all recommendations are expressed in lb/A. These fertilizer rates are based upon typical distances between bed centers for each crop (Table 10). Table 10 also indicates the typical number of planting rows within each bed. Conversions of fertilizer rates from

lb/A to lb/100 LBF, based upon these typical bed spacings, are shown in Table 11.

Use of lb/100 LBF as a fertilizer rate assures that an appropriate rate of fertilizer will be applied, regardless of the total number LBF in the cropped area. In other words, use of lb/A to express the fertilizer rate requires an adjustment based upon actual cropped area.

In reality, the goal is to provide a specific concentration of nutrients to plant roots; that is, a specific amount of fertilizer within a certain volume of soil. This conceptual approach makes sense because most plant roots are confined within the volume of soil comprising the bed, especially under the polyethylene in the full-bed mulch system.

See Table 11 for the conversion of fertilizer rates in lb/A to lb/100 LBF. This table is used correctly by 1) determining the typical bed spacing from Table 10 for the crop; 2) locating the column containing the recommended fertilizer rate in lb/A; and 3) reading down the column until reaching the row containing the typical row spacing. This rate, in lb/100 LBF, should be used even in situations where the grower's bed spacing differs from the typical bed spacing.

**Table 10.** Typical bed spacings for vegetables grown in Florida.

Vegetable	Typical Spacing <sup>1</sup> (ft)	Rows of plants per bed
Broccoli	6	2
Muskmelon	5	1
Cabbage	6	2
Pepper	6	2
Cauliflower	6	2
Summer squash	6	2
Cucumber	6	2
Strawberry	4	2
Eggplant	6	1
Tomato	6	1
Lettuce	4	2
Watermelon	8	1

<sup>1</sup> Spacing from the center of one bed to the center of an adjacent bed.

### IRRIGATION MANAGEMENT

Water management and fertilizer management are linked. Changes in one program will affect the efficiency of the other program. Leaching potential is high for the mobile nutrients such as N and K; therefore, over irrigation can result in movement of these nutrients out of the root zone. This could result in groundwater pollution in the case of N. The goal of water management is to keep the irrigation water and the fertilizer in the root zone. Therefore, growers need knowledge of the root zone of the particular crop so that water and fertilizer inputs can be managed in the root zone throughout the season. For subsurface irrigated crops, water table management is critical. The water table should not fluctuate since this can lead to flushing of nutrients from the root zone. Water tables should be maintained at the proper level to ensure

**Table 11.** Conversion of fertilizer rates in lb/A to lb/100 LBF.

Typical bed spacing (ft)	Recommended fertilizer rate in lb/A (N, P <sub>2</sub> O <sub>5</sub> , or K <sub>2</sub> O)									
	20	40	60	80	100	120	140	160	180	200
	<b>Resulting fertilizer rate in lb/100 LBF (N, P<sub>2</sub>O<sub>5</sub>, or K<sub>2</sub>O)</b>									
3	0.14	0.28	0.41	0.55	0.69	0.83	0.96	1.10	1.24	1.38
4	0.18	0.37	0.55	0.73	0.92	1.10	1.29	1.47	1.65	1.83
5	0.23	0.46	0.69	0.92	1.15	1.38	1.61	1.84	2.07	2.30
6	0.28	0.55	0.83	1.10	1.38	1.65	1.93	2.20	2.48	2.76
8	0.37	0.73	1.10	1.47	1.84	2.20	2.57	2.94	3.31	3.68

optimum moisture in the bed without leading to oversaturation of the root zone and potential losses of nutrients. Water tables can be monitored with a section of PVC pipe sunk in the soil with a calibrated float inside the PVC pipe. The calibrated float can be used to determine the exact level of the water table. Tensiometers placed in the root zone of the crop can be used to provide an indication of soil moisture. If properly maintained, tensiometers are a good tool to monitor soil moisture with the resulting information used to make adjustments in irrigation rates. In most sandy soil situations in Florida, tensiometers placed 6 inches deep in the soil in the root zone should be maintained no wetter than about -8 centibars and no drier than about -12 to -15 centibars.

The exact amount of water to apply to a crop can be gauged by evapotranspiration. Evapotranspiration is estimated by measuring the amount of evaporation from a class A weather bureau evaporation pan. The amount of water that is applied is related to the amount of evaporation.

With increased pressure on growers to conserve water and to minimize the potential for nutrient pollution, it becomes extremely important to learn as much as possible about irrigation management. The key is to tie water applications to a soil moisture measuring device such as a tensiometer and an application amount calculated based on evapotranspiration. If water is properly managed then mobile nutrients such as nitrogen should remain in the root zone to benefit the crop. For more information, see Chapter 8, *Principles and Practices of Irrigation Management for Vegetables*, which is part of this publication.

## PLANT TISSUE ANALYSIS

Analysis of plants for nutrient concentration provides a good tool to monitor nutrient management programs. There are basically two approaches to plant tissue testing: standard laboratory analyses based on dried plant parts; and the newly-developed plant sap testing procedures. Both procedures have value in nutrient management programs for vegetable crops, each having its own advantages and disadvantages.

Standard laboratory analyses can be very accurate and are the most quantitative procedure. However, they can be time consuming for most diagnostic situations in the field. Standard laboratory analysis involves analyzing the most-recently-matured leaf of the plant for an array of nutrients. The resulting analyses are compared against published adequate ranges for that particular crop. Laboratory results that fall outside the adequate range for that nutrient may indicate either a deficiency or possibly a toxicity (especially in the case of micronutrients). The most-recently-matured leaf serves well for routine crop monitoring and diagnostic procedures for most nutrients. However, for the

immobile nutrients such as Ca, B, and certain other micronutrients, younger leaves are generally preferred. For vegetable crops grown in Florida, University of Florida publication SS-VEC-42, *Plant Tissue Analysis and Interpretation for Vegetables in Florida*, should be used as a reference when tissue analysis results are obtained from the laboratory. For each crop in this production recommendations booklet, standard laboratory sufficiency ranges are presented for one crop growth stage. For other crop growth stages, refer to SS-VEC-42.

Several plant sap quick test kits have been calibrated for N and K for several vegetables in Florida (Fig. 2-6). These testing kits analyze fresh plant sap for N and K. Quick test kits offer speed of analysis however, are less quantitative than standard laboratory procedures. However, quick tests are accurate enough and if properly calibrated are a valuable tool for on-the-spot monitoring of plant nutrient status with the goal of making fine adjustments in fertilizer application programs, especially for those involving drip irrigation. Results of research with plant sap kits and their calibration is presented in SS-VEC 42, and Circular 1144, *Plant Petiole Sap-Testing - Guide for Vegetable Crops*.

## DRIP IRRIGATION/FERTIGATION

Drip irrigation has become a very important water management tool for Florida vegetable growers (Fig. 2-7). Approximately 60,000 acres of vegetables are produced with drip irrigation yearly in Florida. Many drip irrigation users have turned to fertigation (applying nutrients through the irrigation tube) to gain better fertilizer management capability. In most situations, N and K are the nutrients injected through the irrigation tube. This is because these nutrients are mobile in Florida soils, and can be leached. Split applications of N and K through irrigation systems offers a means to capture management potential and reduce leaching losses. Other nutrients, such as P and micronutrients, are usually applied to the soil rather than by injection. This is because chemical precipitation can occur with these nutrients and the high calcium carbonate content of our irrigation water in Florida.

Nutrient management through irrigation tubes involves precise scheduling of N and K applications. Application rates are determined by crop growth and resulting nutrient demand. Demand early in the season is small and thus rates of application are small, usually on the order of  $\frac{1}{2}$  to  $\frac{3}{4}$  lb of N or  $K_2O$  per acre per day. As the crop grows, nutrient demand increases rapidly so that for some vegetable crops such as tomato the demand might be as high as 2 lb of N or  $K_2O$  per day. Schedules of N and K application have been developed for most vegetables produced with drip irrigation in Florida. Schedules for these crops are presented in the crop chapters in this book. Much more detailed information on nutrient and water management for

drip irrigation systems can be obtained from Circ. 1181 *Fertilizer Application and Management for Micro (or Drip) Irrigated Vegetables in Florida*.

## SOIL PREPARATION

A well-prepared seed or planting bed is important for uniform stand establishment of vegetable crops. Old crop residues should be plowed down well in advance of crop establishment. A 6- to 8-week period between plowing down of green cover crops and crop establishment is recommended to allow the decay of the refuse. Freshly incorporated plant material promotes high levels of damping-off organisms such as *Pythium* spp. and *Rhizoctonia* spp. Turning under plant refuse well in advance of cropping reduces damping-off disease organisms. Land should be kept disced if necessary to keep new weed cover from developing prior to cropping.

Chisel plowing is beneficial in penetrating and breaking tillage pan layers in fields. If plastic mulch culture is practiced, debris and large undecayed roots will create problems in preparing good beds over which mulch will be applied.

## BEDDING

Fields where seepage irrigation is used or fields prone to flooding should be cropped using raised beds. Beds generally range from 3 to 8 inches in height, with high beds of 6 to 8 inches preferred where risk of flooding is greatest. Raised beds dry faster than if the soil was not bedded, requiring closer attention to irrigation management especially early in the season when root systems are limited. Raised beds promote early season soil warming resulting in somewhat earlier crops during cool seasons. Many raised beds covered with mulch in north Florida in sandy, well drained soils do not need to be as high as 6 to 8 inches as they do in poorly drained soils.

Bedding equipment may include single or double bedding discs, and curved bedding blades. After the soil is cut and thrown into a loose bed the soil is usually firmed with a bed press. In unmulched production the loosely formed bed may be leveled off at the top by dragging a board or bar across the bed top. Boarding-off the raised beds is common in unmulched watermelon production in central and northern Florida. Mulching requires a smooth, well-pressed bed for efficient heat transfer from black mulch to the soil. Adequate soil moisture is essential in forming a good bed for mulching. Dry sandy soils will not form a good bed for a tight mulch application. Overhead irrigation is sometimes needed to supply adequate moisture to dry soils before bedding.

## COVER CROPS

Cover crops between vegetable cropping seasons can provide several benefits. The use of cover crops as green manure can slightly increase soil organic matter during the growing season. Properties of soil tilth can also be improved with turning under good cover crops. The cover can reduce soil losses due to erosion from both wind and water. Many crops are effective at recycling nutrients left from previous crops. Recycling of nutrients is becoming an increasingly important issue in protecting groundwater quality.

The selection of a cover crop is based on the seasonal adaptation and intended use for the crops. Vegetable production in south Florida results in cover crops needed during the late spring and summer months. Summer grasses like sorghum or sudan/sorghum hybrids have been popular among Florida producers as a summer cover. Pearl millet is another grass crop providing excellent cover but is not as popular as sudan/sorghum. Both pearl millet and sudan/sorghum provide a vigorous tall crop with high biomass production and are excellent at competing with weeds. The cover crop selected should have resistance to nematodes or at least serve as a relatively poor nematode host. Warm-season legumes such as velvet bean and hairy indigo have been noted for their resistance to nematodes. Hairy indigo has been unpopular because of its habit of reseeding. It also has hard seed and produces volunteers in later years. Alyceclover is another warm season legume with one variety, F1-3, having nematode resistance. Alyceclover produces an excellent quality hay for producers that can utilize hay from a cover crop.

In north Florida, vegetable crops are established in the spring and early fall. Cover crops are generally utilized during the winter months of November through March. Popular cool season grasses have included rye, wheat, oats, or ryegrass. The traditional crop rotation for watermelon growers has included the use of well-established bahia grass pastures followed by a crop of watermelon. The acreage of available bahia grass pastures for rotation has been reduced and these pastures are difficult to find for many growers. As a result, growers are being forced to more intensively crop fields. Cover crops would be helpful in managing the land. When bahia grass sod is used for production, the extensive root system must be very well tilled well in advance of the cropping season to break up the clumps, especially if plastic mulch will be used. Deep plowing is best to facilitate decomposition of the grass roots and stems.

## WINDBREAKS

The use of windbreaks is an important cultural practice consideration in many vegetable crops and in most states

in the United States. Windbreaks used in agriculture are barriers, either constructed or vegetative, of sufficient height to create a windless zone to their leeward or protected side. Strong winds, even if a few hours in duration, can cause injury to vegetable crops by: whipping plants around, abrasion with solid particles (“sand blasting”), cold damage, and plant dessication. Windbreaks are especially important to protect young plants that are most susceptible to wind damage. Abrasion to plants from wind-blown sand is of concern in most of Florida where sandy soils are commonly used for production. Spring winds in Florida are expected each year. Many of the vegetable crops produced in central and north Florida are at a young and very susceptible stage during these windy spring periods. Strips of planted rye are generally recommended for temporary windbreaks in those areas (Fig. 2-8). Sugarcane can also serve as a more permanent windbreak in South Florida (Fig. 2-9).

The primary reasons windbreaks have been used in vegetable crops has been to reduce the physical damage to the crop from the whipping action of the wind and to reduce sand blasting. Young, unprotected vegetable crops stands can be totally lost from these two actions. Many Florida vegetable crops are grown using plastic mulch culture. Young cucurbit crops, such as watermelon and cantaloupe grown on plastic are especially susceptible to the whipping action of the wind. Vines of these crops eventually become anchored to the soil between mulched beds, however, young vines can be whipped around in circles for several days until they become anchored. The physical damage by whipping and sandblasting can reduce stand, break or weaken plants, open wounds which can increase disease, and reduce flowering and fruit set.

Windbreaks can also help conserve moisture for the crop. Effective windbreaks reduce the wind speed reaching the crops. This reduces both direct evaporation from the soil and transpiration losses from the plant. Improved moisture conditions can help in early season stand establishment and crop growth. Air temperatures around the crop can also be slightly modified by windbreaks. Temperature on the leeward side of the windbreaks can be slightly higher than if no windbreak were present. Early season crop growth is also greater when windbreaks are utilized. Workers in several states reported increased earliness when rye strips were effectively used as windbreaks.

A field layout to include windbreaks must be properly designed to achieve the maximum benefit. The windbreaks should be positioned perpendicular to the prevailing winds. This determination is perhaps more difficult in Florida than most other states, however, windbreaks planned for protection in the spring should generally protect against winds from the west or northwest. Wind protection is achieved as long as the barrier is a least three feet high, the vegetation

is sufficiently dense, and is positioned perpendicular to the prevailing wind.

The height of the windbreak is the most important factor in determining how far apart the strips must be located. Research on windbreaks has been conducted indicating wind protection is afforded to a distance of 6 to 20 times the height of the barrier. Field research with rye strips showed protection was afforded up to a distance of 10 times the height of the barrier. For example, a healthy crop of rye planted in a 5 to 8 ft wide strip using a grain drill and reaching a height of 3 ft would afford wind protection up to 30 ft from the rye strip. If the same rye strip reached a height of 4 ft it would afford protection up to 40 ft from the rye strip. These examples use the calculation of protection afforded up to 10 times the height of an adequate rye strip.

Crops such as small grains, trees, shrubs, or sugarcane are “permeable” barriers in comparison to solid barriers such as smooth constructed walls. Solid barriers are less effective windbreaks than permeable barriers. Wind passing over a solid barrier is deflected over and creates an area of turbulence on the protected side and returns to the ground quickly.

Another type of technology that can provide excellent protection from high winds is the use of plastic row tunnels. Polyethylene or polypropylene materials are placed over the plants in a row and held in place. Tunnels are popular for many vegetable crops, especially cucurbits such as cantaloupes. The cover is removed from cucurbits when the first female blooms appear to allow honeybees to pollinate the crops. Tunnels are generally used in conjunction with rye strips because the tunnels have to be removed and once removed the crop is susceptible to wind.

The most widely used windbreak in vegetable crops across the United States is the rye strip method. Winter or cereal rye (*Secale cereale*) is the preferred small grain for this use because the seed is usually cheaper, it provides more growth under cold temperatures and results in the highest plant habit. In some cases the field is solid seeded and later tilled in only the narrow strips where the plastic mulch bed is applied. This leaves a narrow strip of rye between each bed or row and is generally a very effective windbreak design. This design can result in more difficulties in weed management if weeds emerge in the rye strips, however, the rye can be managed with herbicide in certain crops.

The most common use of rye as a windbreak is planting it into strips. Seeding rye should be done in the fall (October - December) for protection in a spring crop. The strips are typically 5-8 ft wide and planted with a grain drill. The windbreak is a valuable component of the cropping system and should be treated as such. A top dressing

or two of a fertilizer (at least nitrogen) will promote sufficient early spring growth of the rye to maximize effectiveness as a windbreak. Unfertilized rye strips on low fertility soil will often result in poor, thin, short strips of rye that will be less effective as a windbreak.

The spacing of the rye strips every 30 to 40 feet allows them to also be used as drive roads or spray roads in the field. These are generally necessary in managing most vegetable crops and therefore the rye strips are not taking away cropped areas of the field.

When the rye strips have served their purpose, they can be removed by mowing, rototilling, or discing. If mowing is used in a plastic mulched field, the mower should not throw the rye stems into the plastic area because holes will be pierced in the mulch. One insect management concern in using rye strips in Florida is their attractiveness to thrips. Rye strips also seem to be an excellent environment for beneficial insects, especially lady beetles. If thrips need to be managed in the rye strips, the strips could be sprayed just before the rye is mowed or tilled out. Once the rye is destroyed, the thrips migrate to the crops so control would be more effective while they are still on the rye strips.